

### Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Donkey Polyclonal, Peroxidase (Horseradish)
Catalog # ASR1930

### **Specification**

**Physical State** 

Host Isotype

**Immunogen** 

Reconstitution Volume

Reconstitution Buffer

# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-HUMAN IgG (H&L) (DONKEY) Antibody

**Peroxidase Conjugated** 

Host **Donkey** 

Conjugate Peroxidase (Horseradish)

Target Species

Clonality

Application

Human

Polyclonal

WB, E, IC

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:10,000;Immunochemistry

1:500-1:2,500 Lyophilized

IgG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Human IgG whole molecule

1.0 mL

Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information

### **Shipping Condition**

**Ambient** 

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Human IgG and Human Serum.

## **Storage Condition**

Store vial at  $4^{\circ}$  C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at  $4^{\circ}$  C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

## Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images