

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2040

Specification

Physical State

Host Isotype

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-HUMAN IgG IgA IgM (H&L) (GOAT)

Antibody Biotin Conjugated

Host Goat
Conjugate Biotin
Target Species Human
Clonality Polyclonal
Application WB, IHC, E

Application Note ELISA 1:400,000; Western Blot

1:5,000-1:20,000;Immunohistochemistry

1:1,000-1:5,000 Lyophilized

IaG

Target Isotype IgG IgA IgM

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Human IgG, IgA and IgM whole molecules

1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Reconstitution Volume

Ambient

Purity

This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin and anti-Goat Serum. This product is suitable for the detection of all Human immuno-globulin classes, isotypes and chain combinations.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Images