

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody Donkey Polyclonal, Fluorescein (FITC)

Catalog # ASR2079

Specification

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description	F(ab')2 Anti-GOAT IgG [H&L] (DONKEY) Antibody Fluorescein Conjugated Min X Ch GP Ham Hs Hu Ms Rb & Rt Serum Proteins
Host	Donkey
Conjugate	Fluorescein (FITC)
FP Value	5. moles Fluorescein (FITC) per mole of lgG F(ab')2
Target Species	Goat
Clonality	Polyclonal
Application	IF, FC
Application Note	FLISA 1:10,000-1:50,000;IF Microscopy
	1:1,000-1:5,000;FlowCytometry
	1:500-1:2,500
Physical State	Lyophilized
Host Isotype	lgG F(ab')2
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M
Immunagan	Sodium Chloride, pH 7.2
Immunogen Reconstitution Volume	Goat IgG whole molecule 500 μL
Reconstitution Buffer	Restore with deionized water (or
Reconstitution bullet	equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C



or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody - Images

F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody -Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.