

Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody

Rabbit Polyclonal, Biotin Catalog # ASR2156

Specification

Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-DOG IgG F(ab')2 (RABBIT)

Antibody Biotin Conjugated

Host Rabbit
Conjugate Biotin
Target Species Dog

Clonality Polyclonal Application WB, IHC, E

Application Note ELISA 1:20,000-1:100,000;Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000 Lyophilized

Physical State
Host Isotype
IgG

Target Isotype IgG F(ab')2

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Dog IgG F(ab')2 fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Dog IgG, Dog IgG F(ab')2 and Dog Serum. No reaction was observed against Dog IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody - Protein Information



Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Dog IgG F(ab')2 (Biotin Conjugated) Secondary Antibody - Images