

Anti-Mouse IgG F(c) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2198

Specification

Anti-Mouse IgG F(c) Secondary Antibody - Product Information

Description Anti-MOUSE IgG F(c) (GOAT) Antibody

Biotin Conjugated

Host Goat Biotin Conjugate **Target Species** Mouse Reactivity Mouse Clonality **Polyclonal** Application WB, E, IC

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000; Immunochemistry

1:1.000-1:5.000 Lvophilized

Physical State Host Isotype laG Target Isotype IaG F(c)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Anti-mouse IgG F(c) was produced by

repeated immunization with mouse IgG

F(c) fragment in goat.

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Thimerosal

Anti-Mouse IgG F(c) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG F(c) Secondary Antibody - Protein Information

Anti-Mouse IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Mouse IgG F(c) Secondary Antibody - Images

Anti-Mouse IgG F(c) Secondary Antibody - Background

Anti-Mouse IgG F(c) Antibody generated in goat detects specifically mouse IgG F(c). Anti-Mouse IgG F(c) antibody is ideal for investigators involved in Serum Protein Component research. Anti-Mouse IgG F(c) Antibody is ideal for investigators in Cancer, Immunology, and Microbiology research.