

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody
Rabbit Polyclonal, Rhodamine (TRITC)
Catalog # ASR2319**Specification****Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information**

Description	Anti-CAT IgG F(c) (RABBIT) Antibody
Host	Rhodamine Conjugated
Conjugate	Rabbit
FP Value	Rhodamine (TRITC)
Target Species	1.6 moles Rhodamine (TRITC) per mole of IgG
Clonality	Cat
Application	Polyclonal
Application Note	IF, FC
Physical State	FLISA 1:10,000-1:50,000;IF Microscopy
Host Isotype	1:1,000-1:5,000;FlowCytometry
Target Isotype	1:500-1:2,500
Buffer	Lyophilized
Immunogen	IgG
Reconstitution Volume	IgG F(c)
Reconstitution Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer	Cat IgG F(c) fragment
Preservative	1.0 mL
	Restore with deionized water (or equivalent)
	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
	0.01% (w/v) Sodium Azide

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Cat IgG, Cat IgG F(c) and Cat Serum. No reaction was observed against Cat IgG F(ab').

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Cat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.