

# Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2335

### **Specification**

## Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-HORSE IgG F(c) (GOAT) Antibody

**Biotin Conjugated** 

Host Goat
Conjugate Biotin
Target Species Horse
Clonality Polyclonal
Application WB, E, IC

Application Note ELISA 1:20,000-1:100,000;Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized

Physical State
Host Isotype
Target Isotype
Lyophiliz
IgG
IgG F(c)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Horse IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Thimerosal

### Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody - Additional Information

## **Shipping Condition**

**Ambient** 

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Horse IgG, Horse IgG F(c) and Horse Serum. No reaction was observed against Horse IgG F(ab')2.

### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



## Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody - Protein Information

## Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Horse IgG F(c) (Biotin Conjugated) Secondary Antibody - Images