

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal. Peroxidase (Horseradish) Catalog # ASR2453

Specification

Physical State

Host Isotype

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Product **Information**

Description Anti-GUINEA PIG IgG F(ab')2 (GOAT)

Antibody Peroxidase Conjugated

Host Goat

Peroxidase (Horseradish) Conjugate

Target Species Guinea Pig Clonality **Polyclonal** Application WB, E, IC

Application Note ELISA 1:2,000-1:6,000; Western Blot

1:1,000-1:10,000; Immunochemistry

1:500-1:2,500 Lyophilized

laG

Target Isotype IgG F(ab')2

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Guinea Pig IgG F(ab')2 fragment **Immunogen** Reconstitution Volume

1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Additional **Information**

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Guinea Pig IgG, Guinea Pig IgG F(ab')2 and Guinea Pig Serum. No reaction was observed against Guinea Pig IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Guinea Pig IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Images