

#### Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2470

#### **Specification**

#### Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-HUMAN IgG F(c) (RABBIT) Antibody

**Peroxidase Conjugated** 

Host Rabbit

Conjugate Peroxidase (Horseradish)

Target Species
Clonality
Application
Human
Polyclonal
WB, E, IC

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:5,000;Immunochemistry

1:500-1:2,500 Lyophilized

Physical State
Host Isotype
Target Isotype
IgG F(c)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Human IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

# Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information

# **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')2

## **Storage Condition**

Store vial at  $4^{\circ}$  C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at  $4^{\circ}$  C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



# Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information

# Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images