

**Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody**  
**Goat Polyclonal, Peroxidase (Horseradish)**  
**Catalog # ASR2488****Specification****Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Product Information**

|                       |   |
|-----------------------|---|
| Description           | <b>Anti-RABBIT IgG F(ab')<sub>2</sub> (GOAT) Antibody Peroxidase Conjugated (Min X Human Serum Proteins)</b>                                  |
| Host                  | <b>Goat</b>   |
| Conjugate             | <b>Peroxidase (Horseradish)</b>   |
| Target Species        | <b>Rabbit</b>   |
| Reactivity            | <b>Rabbit</b>   |
| Clonality             | <b>Polyclonal</b>   |
| Application           | <b>WB, E, IC</b>  |
| Application Note      | <b>ELISA 1:130,000;Western Blot 1:2,000-1:20,000;Immunochemistry 1:1,000-1:10,000</b>   |
| Physical State        | <b>Lyophilized</b>  |
| Host Isotype          | <b>IgG</b>  |
| Target Isotype        | <b>IgG F(ab')<sub>2</sub></b>   |
| Buffer                | <b>0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>  |
| Immunogen             | <b>Anti-Rabbit IgG F(ab')<sub>2</sub> fragment was produced by repeated immunization with Rabbit IgG F(ab')<sub>2</sub> fragment in goat.</b> |
| Reconstitution Volume | <b>1.0 mL</b>   |
| Reconstitution Buffer | <b>Restore with deionized water (or equivalent)</b>   |
| Stabilizer            | <b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>   |
| Preservative          | <b>0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!</b>   |

**Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

Anti-RABBIT IgG F(ab')<sub>2</sub> (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab')<sub>2</sub> and Rabbit Serum. No reaction was observed against Rabbit IgG F(c) or Human Serum Proteins.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.


### **Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**

### **Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images**



Western Blot of Anti-Rabbit IgG F(ab')<sub>2</sub> (GOAT) Antibody . Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule . Lane 2: Rabbit IgG F(ab) Fragment . Lane 3: Rabbit IgG F(c) Fragment . Lane 4: Rabbit IgM Whole Molecule . Lane 5: Normal Rabbit Serum . All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG F(ab')<sub>2</sub> (GOAT) Antibody 1:10,000 for 60 min at RT. Secondary antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in MB-070 for 30 min at RT. Predicted/Observed Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.

### **Anti-Rabbit IgG F(ab')<sub>2</sub> (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Background**

Anti-Rabbit IgG F(ab')<sub>2</sub> fragment antibody generated in goat detects specifically Rabbit IgG F(ab')<sub>2</sub> fragment. This secondary antibody anti-Rabbit is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.