

## F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody Goat Polyclonal, Biotin

Catalog # ASR2518

## Specification

# F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note	F(ab')2 Anti-GUINEA PIG IgG (H&L) (GOAT) Antibody Biotin Conjugated Goat Biotin Guinea Pig Polyclonal WB, IHC, E ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	lgG F(ab')2
Target Isotype	lgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M
	Sodium Chloride, pH 7.2
Immunogen	Guinea Pig IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or
	equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) -
	Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

# F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

#### Shipping Condition Ambient

#### Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

#### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



#### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

# F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

### F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images

## F(ab')2 Anti-Guinea Pig IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.