

Anti-Goat IgG (H&L) Secondary Antibody

Rabbit Polyclonal, Biotin Catalog # ASR2626

Specification

Physical State

Host Isotype

Anti-Goat IgG (H&L) Secondary Antibody - Product Information

Anti-GOAT IgG (H&L) (RABBIT) Antibody Description

Biotin Conjugated (Min X Human Serum

Proteins) **Rabbit Biotin**

Host Conjugate **Target Species** Goat Clonality **Polyclonal** Application WB, E, IC

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000; Immunochemistry

1:1.000-1:5.000 Lvophilized

laG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Goat IgG whole molecule

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Goat IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

Anti-Goat IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against Human Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Goat IgG (H&L) Secondary Antibody - Protein Information

Anti-Goat IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Goat IgG (H&L) Secondary Antibody - Images

Anti-Goat IgG (H&L) Secondary Antibody - Background

Anti-Goat IgG (H&L) Antibody generated in rabbit detects specifically goat IgG (H&L). This secondary Anti-Goat IgG (H&L) Antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. Anti-Goat IgG (H&L) Antibody is ideal for researchers in Immunology, Cancer, and Microbiology research.