

## Anti-MOUSE IgG2a (Rhodamine Conjugated) Pre-adsorbed Secondary Antibody

**Goat Polyclonal, Rhodamine (TRITC)** Catalog # ASR2708

### **Specification**

## Anti-MOUSE IgG2a (Rhodamine Conjugated) Pre-adsorbed Secondary Antibody - Product **Information**

Description Anti-MOUSE IgG2a (Gamma 2a chain)

> (GOAT) Antibody Rhodamine Conjugated (Min Cross Bv, Hu, and Rb Serum Proteins)

Goat

Host **Rhodamine (TRITC)** Conjugate

FP Value 3.1 moles Rhodamine (TRITC) per mole of

IqG

**Target Species** Mouse Reactivity Mouse Clonality **Polyclonal Application** WB, IF, IC

**Application Note** FLISA 1:20,000-1:100,000; IF Microscopy

1:1,000-1:5,000; Western Blot

1:2,000-1:10,000; Immunochemistry

1:1,000-1:5,000 Lyophilized

**Physical State** Host Isotype **IgG** Target Isotype IqG2a

**Buffer** 0.02 M Potassium Phosphate, 0.15 M

> Sodium Chloride, pH 7.2 Mouse IgG2a heavy chain

Immunogen Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

## Anti-MOUSE IgG2a (Rhodamine Conjugated) Pre-adsorbed Secondary Antibody -**Additional Information**

**Shipping Condition** 

**Ambient** 

### **Purity**

MOUSE IgG2a Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse Serum and Mouse IgG2a. Specificity was confirmed by ELISA at less than 1% cross-reactivity against other mouse heavy or light chain isotypes. No reaction was observed against Bovine, Human, and Rabbit Serum Proteins. Specificity was confirmed by ELISA at less than 1% of target signal.



## **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.


## Anti-MOUSE IgG2a (Rhodamine Conjugated) Pre-adsorbed Secondary Antibody - Protein Information

## Anti-MOUSE IgG2a (Rhodamine Conjugated) Pre-adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


#### Anti-MOUSE IgG2a ( Rhodamine Conjugated) Pre-adsorbed Secondary Antibody - Images



Dot Blot of Goat anti-Mouse IgG2a Antibody Rhodamine Conjugated Pre-absorbed. Antigen: Mouse IgG2a. Load: Lane 1 - 200 ng Lane 2 - 66.7 ng Lane 3 - 22.2 ng Lane 4 - 7.41 ng Lane 5 - 2.47 ng. Primary antibody: n/a. Secondary antibody: Goat anti-Mouse IgG2a Antibody Rhodamine Conjugated Pre-absorbed at 1:1,000 overnight at 4°C. Block: MB-070 for 1 HR at RT.

# Anti-MOUSE IgG2a ( Rhodamine Conjugated) Pre-adsorbed Secondary Antibody - Background

Mouse IgG2a secondary antibody is available in a variety of formats. Rhodamine secondaaries are designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor





imaging, utilizing various commercial platforms.