

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody

Goat Polyclonal, Fluorescein (FITC)
Catalog # ASR2712

Specification

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Product Information

Description Anti-MOUSE IgG1 (Gamma 1 chain) (GOAT)

Antibody Fluorescein Conjugated (Min Cross Bv, Hu, and Rb Serum Proteins)

Host Goat

Conjugate Fluorescein (FITC)

FP Value 2.8 moles Fluorescein (FITC) per mole of

IgG Mor

Target Species Mouse
Reactivity Mouse
Clonality Polyclonal
Application WB, IF, FC, IC

Application Note FLISA 1:10,000-1:50,000;IF Microscopy

1:1,000-1:5,000;FlowCytometry 1:200-1:500;Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized

Physical State Lyop
Host Isotype IgG
Target Isotype IgG1

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Mouse IgG heavy chain

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

Immunogen

Anti-Mouse IgG1 FITC conjugated secondary antibody prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG1 coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Human, and Rabbit Serum Proteins. Specificity was confirmed by ELISA at less than 1% of target signal.



Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Protein Information

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Images



Western Blot of Fluorescein conjugated Goat anti-Mouse IgG1 (Gamma 1 chain) (Pre-Adsorbed) secondary antibody. Lane 1: Mouse IgG1. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Fluorescein goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 55 kDa, 55 kDa for Mouse IgG1 (Gamma 1 chain). Other band(s): none.

Anti-Mouse IgG1 (Fluorescein Conjugated) Pre-adsorbed Secondary Antibody - Background

FITC Conjugated Secondary Antibodies are designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.