

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2824

Specification

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description Anti-Rat IgG IgA IgM (H&L) (GOAT)

Antibody Biotin Conjugated

Host Goat
Conjugate Biotin
Target Species Rat
Reactivity Rat

Clonality Polyclonal Application WB, E, IC

Application Note ELISA 1:150,000; Western Blot

1:2,000-1:20,000;Immunochemistry

1:1,000-1:10,000

Physical State
Host Isotype
Lyophilized
IgG

Target Isotype IgG IgA IgM

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Anti-Rat IgG IgA IgM (H&L) was produced

by repeated immunization with Rat IgG, IgA and IgM whole molecules in goat.

Reconstitution Volume 2.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Rat IgA, Rat IgG and Rat IgM. This reagent is suitable for the detection of all rat immunoglobulin subclasses, isotypes and chain combinations.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted



Tel: 858.875.1900 Fax: 858.875.1999

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Images

Anti-Rat IgG IgA IgM (H&L) (Biotin Conjugated) Secondary Antibody - Background

Anti-Rat IgG IgA IgM (H&L) biotin conjugated antibody generated in goat detects specifically rat IgG IgA IgM (H&L). This secondary biotin conjugated antibody anti-Rat is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.