

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody Rabbit Polyclonal, Texas Red[®] Catalog # ASR2852

Specification

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Product Information

Description Host	Anti-MOUSE IgM (mu chain) (RABBIT) Antibody Texas Red™ Conjugated Rabbit
Conjugate	Texas Red®
FP Value	1.8 moles Texas Red® per mole of lgG
Target Species	Mouse
Clonality	Polyclonal
Application	IF, FC
Application Note	FLISA 1:10,000-1:50,000;IF Microscopy
	1:1,000-1:5,000;FlowCytometry
	1:500-1:2,500
Physical State	Lyophilized
Host Isotype	lgG
Target Isotype	lgM μ chain
Buffer	0.02 M Potassium Phosphate, 0.15 M
	Sodium Chloride, pH 7.2
Immunogen	Mouse IgM mu heavy chain
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Mouse IgM and Mouse Serum. No reaction was observed against Mouse IgG or Mouse IgA.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Protein Information

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Images



Dot Blot of Rabbit anti-Mouse IgM (mu chain) Antibody Texas Red Conjugated. Antigen: Mouse IgM. Load: Lane 1 - 200 ng Lane 2 - 66.7 ng Lane 3 - 22.2 ng Lane 4 - 7.41 ng Lane 5 - 2.47 ng. Primary antibody: n/a. Secondary antibody: Rabbit anti-Mouse IgM (mu chain) Antibody Texas Red Conjugated at 1:1,000 for 60 min at RT. Block: MB-070 for 60 min at RT.

Anti-Mouse IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.