

**F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody**  
**Goat Polyclonal, Texas Red®**  
**Catalog # ASR2868**

### Specification

**F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-HUMAN IgG F(ab')<sub>2</sub> (GOAT) Antibody Texas Red™ Conjugated Min X Bv Hs Ms &amp; Rt Serum Proteins Goat Texas Red® 2.7 moles Texas Red® per mole of IgG F(ab')<sub>2</sub> Human Polyclonal IF, FC FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500 Lyophilized IgG F(ab')<sub>2</sub> IgG F(ab')<sub>2</sub> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Human IgG F(ab')<sub>2</sub> fragment 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide</b>
Host	
Conjugate	
FP Value	
Target Species	
Clonality	
Application	
Application Note	
Physical State	
Host Isotype	
Target Isotype	
Buffer	
Immunogen	
Reconstitution Volume	
Reconstitution Buffer	
Stabilizer	
Preservative	

**F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information**

### Shipping Condition

Ambient

### Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Human IgG, Human IgG F(ab')<sub>2</sub> and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Human IgG F(c) or Bovine, Horse, Mouse, Rabbit, Rat and Sheep Serum Proteins.

### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information****F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Images****F(ab')<sub>2</sub> Anti-Human IgG F(ab')<sub>2</sub> (Texas Red™ Conjugated) Pre-Adsorbed Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.