

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2883

Specification

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description F(ab) Anti-HUMAN IgG (H&L) (GOAT)

Antibody Peroxidase Conjugated

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species

Clonality

Application

Human

Polyclonal

WB, IHC, E

Application Note ELISA 1:10,000-50,000;Western Blot

1:1,000-1:5,000;Immunohistochemistry

Physical State
Host Isotype
Target Isotype

1:500-1:2,500
Lyophilized
IgG F(ab)
IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Human IgG whole molecule

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Immunogen

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Fab Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images