

Fab Anti-Rabbit IgG (H&L) Secondary Antibody
Donkey Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2886**Specification****Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Product Information**

Description	F(ab) Anti-RABBIT IgG (H&L) (DONKEY) Antibody Peroxidase Conjugated
Host	Donkey
Conjugate	Peroxidase (Horseradish)
Target Species	Rabbit
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:8,000-1:32,000;Western Blot 1:1,000-1:3,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG F(ab)
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Rabbit IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

F(ab) Anti-Rabbit IgG (H&L) was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Donkey IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Protein Information

Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Images

Fab Anti-Rabbit IgG (H&L) Secondary Antibody - Background

F(ab) Anti-Rabbit IgG (H&L) antibody generated in Donkey detects specifically Rabbit IgG whole molecule. This F(ab) Anti-Rabbit IgG Secondary Antibody is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays. F(ab) Anti-Rabbit IgG (H&L) Antibody is ideal for investigators in Immunology, Cancer, and Microbiology research.