

Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody
Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3011**Specification****Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	Anti-FERRET IgG IgA IgM (H&L) (GOAT) Antibody Peroxidase Conjugated
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Ferret
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:5,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG IgA IgM
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Ferret IgG IgA and IgM whole molecule was produced by repeated immunization with Ferret IgG IgA and IgM whole molecules in goat.
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Thimerosal

Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition****Ambient****Purity**

This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Ferret IgG, Ferret IgA and Ferret IgM. This reagent is suitable for the detection of all Ferret immunoglobulin subclasses and chain combinations.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information**Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Images**Anti-Ferret IgG IgA IgM (H&L) (Peroxidase Conjugated) Secondary Antibody - Background**

Anti-Ferret IgG IgA and IgM whole molecule antibody generated in goat detects specifically Ferret IgG IgA and IgM whole molecules. This secondary antibody anti-Ferret is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.