

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR3340

Specification

Physical State

Host Isotype

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Product **Information**

Description Anti-BOVINE IgG F(ab')2 (GOAT)

Antibody Peroxidase Conjugated

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species Bovine Clonality **Polyclonal** Application WB, IHC, E

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:10,000;Immunohistochemistry

1:500-1:2,500 Lyophilized

laG

Target Isotype IgG F(ab')2

0.01 M Sodium Phosphate, 0.15 M Sodium Buffer

Chloride, pH 7.2

Immunogen Bovine IgG F(ab')2 fragment Reconstitution Volume

2.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Additional **Information**

Shipping Condition

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Bovine IgG, Bovine IgG F(ab')2 and Bovine Serum. No reaction was observed against Bovine IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.



Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Bovine IgG F(ab')2 (Peroxidase Conjugated) Secondary Antibody - Images