

# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR3384

## **Specification**

**Physical State** 

Host Isotype

## Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product **Information**

Description Anti-HUMAN IgG (H&L) (GOAT) Antibody

**Peroxidase Conjugated** 

Goat Host

Conjugate Peroxidase (Horseradish)

**Target Species** Human Clonality **Polyclonal** Application WB, IHC, E

**Application Note** ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:10,000;Immunohistochemistry

1:500-1:2,500 Lyophilized

laG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

> Sodium Chloride, pH 7.2 Human IgG whole molecule

**Immunogen** 2.0 mL

Reconstitution Volume

Reconstitution Buffer Restore with deionized water (or

equivalent)

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional **Information**

#### **Shipping Condition**

**Ambient** 

#### **Purity**

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum.

## **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

# **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

# Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images