

**F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody**  
**Rabbit Polyclonal, Peroxidase (Horseradish)**  
**Catalog # ASR3441****Specification****F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-DOG IgG (H&amp;L) (RABBIT) Antibody Peroxidase Conjugated</b>
Host	<b>Rabbit</b>
Conjugate	<b>Peroxidase (Horseradish)</b>
Target Species	<b>Dog</b>
Clonality	<b>Polyclonal</b>
Application	<b>WB, IHC, E</b>
Application Note	<b>ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:10,000;Immunohistochemistry 1:500-1:2,500</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab')<sub>2</sub></b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Dog IgG whole molecule</b>
Reconstitution Volume	<b>2.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>
Preservative	<b>None</b>

**F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

F(ab')<sub>2</sub> Anti-DOG IgG Antibody is Peroxidase Conjugated. The F(ab')<sub>2</sub> fragment of a rabbit IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Dog IgG and Dog Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti-Pepsin.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information****F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images****F(ab')<sub>2</sub> Anti-Dog IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Background**

Anti-DOG IgG Antibody is a F(ab')<sub>2</sub> fragment and is Peroxidase Conjugated. Anti-Dog antibody detects Dog or canine IgG heavy and light chains. F(ab')<sub>2</sub> Anti-Dog antibody is ideal for investigators involved in serum protein component research.