

# Anti-Rabbit IgG F(c) Secondary Antibody

Donkey Polyclonal, Unconjugated Catalog # ASR3540

### Specification

# Anti-Rabbit IgG F(c) Secondary Antibody - Product Information

Description Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Preservative

Anti-RABBIT IgG F(c) (DONKEY) Antibody Donkey Unconjugated Rabbit Polyclonal WB, E, IC ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Lyophilized laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Rabbit IgG F(c) fragment 5.0 mL Restore with deionized water (or equivalent) 0.01% (w/v) Sodium Azide

## Anti-Rabbit IgG F(c) Secondary Antibody - Additional Information

Shipping Condition Ambient

#### **Purity**

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab')2.

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Rabbit IgG F(c) Secondary Antibody - Protein Information



# Anti-Rabbit IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Rabbit IgG F(c) Secondary Antibody - Images