

## Anti-Turkey IgG (H&L) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR3548

### **Specification**

## Anti-Turkey IgG (H&L) Secondary Antibody - Product Information

Anti-TURKEY IgG (H&L) (RABBIT) Antibody Description

Host Rabbit

Conjugate Unconjugated **Target Species Turkey** Clonality **Polyclonal Application** WB, IHC, E

**Application Note** ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000 Lyophilized

**Physical State** laG

Host Isotype

Target Isotype IaG (H&L)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Turkey IgG whole molecule **Immunogen** 

Reconstitution Volume

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

## Anti-Turkey IgG (H&L) Secondary Antibody - Additional Information

## **Shipping Condition**

**Ambient** 

#### **Purity**

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Turkey IgG and Turkey Serum.


#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

## **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Turkey IgG (H&L) Secondary Antibody - Protein Information





# **Anti-Turkey IgG (H&L) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Turkey IgG (H&L) Secondary Antibody - Images