

RABBIT IgG (BULK ORDER)
Catalog # ASR3580**Specification**

RABBIT IgG (BULK ORDER) - Product Information

| | |
|-----------------------|---|
| Description | RABBIT IgG whole molecule (BULK ORDER) |
| Conjugate | Unconjugated |
| Physical State | Lyophilized |
| Host Isotype | IgG |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Species of Origin | Rabbit |
| Reconstitution Volume | 5.0 mL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Preservative | 0.01% (w/v) Sodium Azide |

RABBIT IgG (BULK ORDER) - Additional Information**Shipping Condition**

Ambient

Purity

Rabbit IgG whole molecular was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Rabbit IgG whole molecular was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit IgG and anti-Rabbit Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Rabbit IgG whole molecule is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


RABBIT IgG (BULK ORDER) - Protein Information**RABBIT IgG (BULK ORDER) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RABBIT IgG (BULK ORDER) - Images



SDS-Page of Rabbit IgG. Lane 1: Rabbit IgG - Non-Reduced. Lane 2: Rabbit IgG - Reduced. Load: 1.0 ug per lane. Predicted/Observed Size: Non-reduced - 130 kDa, Reduced - 55 and 28 kDa for Rabbit IgG. Other Band(s): None.



SDS-PAGE of Rabbit IgG Whole Molecule. Lane 1: Non-reduced Rabbit IgG Whole Molecule. Lane 2: 5 µL OPAL Pre-stained Marker (MB-210-0500). Lane 3: Reduced Rabbit IgG Whole Molecule. Load: 1 µg per lane. Predicted/Observed size: Non-reduced at 150-170 kDa , Reduced at 55, 25 kDa.

RABBIT IgG (BULK ORDER) - Background

Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present.