

Anti-ATDC (RABBIT) Antibody
ATDC Antibody
Catalog # ASR5463**Specification**

Anti-ATDC (RABBIT) Antibody - Product Information

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	This affinity purified antibody has been tested for use in ELISA and western blotting. Specific conditions for reactivity and detection of ATDC should be optimized by the end user. Expect a band approximately ~66 kDa in size corresponding to ATDC by western blotting in the appropriate cell lysate or extract.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to an internal portion of human ATDC protein around lysine 116.
Preservative	0.01% (w/v) Sodium Azide

Anti-ATDC (RABBIT) Antibody - Additional Information**Gene ID** 23650**Other Names**
23650**Purity**

This product was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with over-expressed, acetylated and non-acetylated (at K116) ATDC protein. A BLAST analysis was used to suggest cross-reactivity with ATDC from human, horse, bovine, chimpanzee and macaque based on a 100% homology with the immunizing sequence. Partial reactivity is expected against rat and mouse ATDC based on 92% homology with the immunizing sequence. Cross-reactivity with ATDC from other sources has not been determined.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-ATDC (RABBIT) Antibody - Protein Information

Name TRIM29

Synonyms ATDC

Function

Plays a crucial role in the regulation of macrophage activation in response to viral or bacterial infections within the respiratory tract. Mechanistically, TRIM29 interacts with IKBKG/NEMO in the lysosome where it induces its 'Lys-48' ubiquitination and subsequent degradation. In turn, the expression of type I interferons and the production of pro-inflammatory cytokines are inhibited. Additionally, induces the 'Lys-48' ubiquitination of STING1 in a similar way, leading to its degradation.

Cellular Location

Cytoplasm. Lysosome. Note=Colocalizes with intermediate filaments

Tissue Location


Expressed in placenta, prostate and thymus.

Anti-ATDC (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ATDC (RABBIT) Antibody - Images



Rockland's affinity purified anti-ATDC antibody shows reactivity by dot blot with acetylated and non-acetylated forms of the immunizing peptide. This antibody is predicted to recognize both acetylated (AcK116) and non-acetylated forms of ATDC protein. Personal communication, Z. Yuan, H Lee Moffitt Cancer Center and Research Institute.

Anti-ATDC (RABBIT) Antibody - Background

Ataxia-telangiectasia group D-associated protein (ATDC), also called tripartite motif-containing protein 29 (TRIM29), is a novel Histone deacetylase (HDAC) associated protein. Its function is tightly regulated by HDAC. ATDC Lysine 116 (K116) is acetylated and has a significant functional role in regulating cell survival and tumorigenesis. ATDC is expressed in placenta, prostate and thymus, and is over expressed in pancreatic and cervical tumors. Its function in tumor cells is not fully understood. It is constitutively phosphorylated by PKC on serine/threonine in A431 cells. The ATDC gene product is one of a group of proteins that share multiple zinc finger motifs and an adjacent leucine zipper motif. These proteins have been proposed to form homo- or heterodimers involved in nucleic acid binding, consistent with the fact that many of these proteins appear to be transcriptional regulatory factors involved in carcinogenesis and/or differentiation. The likelihood that the ATDC gene product is involved in transcriptional regulation could explain the pleiomorphic characteristics of AT, including abnormal cell cycle regulation.