

## EXOC4 Antibody (monoclonal) (M06)

Mouse monoclonal antibody raised against a partial recombinant EXOC4. Catalog # AT1964a

#### Specification

## EXOC4 Antibody (monoclonal) (M06) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW

WB, E <u>Q96A65</u> <u>NM\_021807</u> Human, Mouse, Rat mouse Monoclonal IgG2a Kappa 110498

### EXOC4 Antibody (monoclonal) (M06) - Additional Information

Gene ID 60412

**Other Names** Exocyst complex component 4, Exocyst complex component Sec8, EXOC4, KIAA1699, SEC8, SEC8L1

**Target/Specificity** EXOC4 (NP\_068579, 1 a.a. ~ 109 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** EXOC4 Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.

### EXOC4 Antibody (monoclonal) (M06) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


EXOC4 Antibody (monoclonal) (M06) - Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (37.73 KDa).




EXOC4 monoclonal antibody (M06), clone 4F1. Western Blot analysis of EXOC4 expression in PC-12((Cat # AT1964a )







EXOC4 monoclonal antibody (M06), clone 4F1 Western Blot analysis of EXOC4 expression in Hela S3 NE ( (Cat # AT1964a )



EXOC4 monoclonal antibody (M06), clone 4F1. Western Blot analysis of EXOC4 expression in Raw 264.7((Cat # AT1964a )



EXOC4 monoclonal antibody (M06), clone 4F1. Western Blot analysis of EXOC4 expression in NIH/3T3((Cat # AT1964a )



Detection limit for recombinant GST tagged EXOC4 is approximately 0.1ng/ml as a capture antibody.

# EXOC4 Antibody (monoclonal) (M06) - Background

The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling



and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

## EXOC4 Antibody (monoclonal) (M06) - References

1.Exocyst subunits are involved in isoproterenol-induced amylase release from rat parotid acinar cells.Imai A, Yoshie S, Haga-Tsujimura M, Nashida T, Shimomura H.Eur J Oral Sci. 2012 Apr;120(2):123-31. doi: 10.1111/j.1600-0722.2012.00952.x.