

HERPUD1 Antibody (monoclonal) (M04)

Mouse monoclonal antibody raised against a partial recombinant HERPUD1. Catalog # AT2354a

Specification

HERPUD1 Antibody (monoclonal) (M04) - Product Information

Application WB, IHC, IP, E **Primary Accession** 015011 Other Accession NM 014685 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 43720

HERPUD1 Antibody (monoclonal) (M04) - Additional Information

Gene ID 9709

Other Names

Homocysteine-responsive endoplasmic reticulum-resident ubiquitin-like domain member 1 protein, Methyl methanesulfonate (MMF)-inducible fragment protein 1, HERPUD1, HERP, KIAA0025, MIF1

Target/Specificity

HERPUD1 (NP_055500, 74 a.a. \sim 180 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 IHC~~1:100~500 IP~~N/A E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

HERPUD1 Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.


HERPUD1 Antibody (monoclonal) (M04) - Protocols

Provided below are standard protocols that you may find useful for product applications.




- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HERPUD1 Antibody (monoclonal) (M04) - Images




Immunoprecipitation of HERPUD1 transfected lysate using anti-HERPUD1 monoclonal antibody and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with HERPUD1 MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.51 KDa).






HERPUD1 monoclonal antibody (M04), clone 2G7 Western Blot analysis of HERPUD1 expression in HepG2 ((Cat # AT2354a)



Western Blot analysis of HERPUD1 expression in transfected 293T cell line by HERPUD1 monoclonal antibody (M04), clone 2G7.


Lane 1: HERPUD1 transfected lysate(44 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to HERPUD1 on formalin-fixed paraffin-embedded human tonsil. [antibody concentration 3 ug/ml]





Detection limit for recombinant GST tagged HERPUD1 is approximately 0.1ng/ml as a capture antibody.

HERPUD1 Antibody (monoclonal) (M04) - Background

The accumulation of unfolded proteins in the endoplasmic reticulum (ER) triggers the ER stress response. This response includes the inhibition of translation to prevent further accumulation of unfolded proteins, the increased expression of proteins involved in polypeptide folding, known as the unfolded protein response (UPR), and the destruction of misfolded proteins by the ER-associated protein degradation (ERAD) system. This gene may play a role in both UPR and ERAD. Its expression is induced by UPR and it has an ER stress response element in its promoter region while the encoded protein has an N-terminal ubiquitin-like domain which may interact with the ERAD system. This protein has been shown to interact with presentlin proteins and to increase the level of amyloid-beta protein following its overexpression. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms. The full-length nature of all transcript variants has not been determined.

HERPUD1 Antibody (monoclonal) (M04) - References

1.BRSK2 is regulated by ER stress in protein level and involved in ER stress-induced apoptosis. Wang Y, Wan B, Li D, Zhou J, Li R, Bai M, Chen F, Yu L.Biochem Biophys Res Commun. 2012 Jun 16.2. Decreased ER-associated degradation of ?-TCR induced by Grp78 depletion with the SubAB cytotoxin. Lass A, Kujawa M, McConnell E, Paton AW, Paton JC, Wojcik C.Int J Biochem Cell Biol. 2008;40(12):2865-79. Epub 2008 Jun 20.