

**MCP Antibody (monoclonal) (M07)****Mouse monoclonal antibody raised against a partial recombinant MCP.****Catalog # AT2822a****Specification**

---

**MCP Antibody (monoclonal) (M07) - Product Information**

Application	<b>E</b>
Primary Accession	<a href="#">P15529</a>
Other Accession	<a href="#">BC030594</a>
Reactivity	<b>Human</b>
Host	<b>mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG2a Kappa</b>
Calculated MW	<b>43747</b>

**MCP Antibody (monoclonal) (M07) - Additional Information****Gene ID** 4179**Other Names**

Membrane cofactor protein, TLX, Trophoblast leukocyte common antigen, CD46, CD46, MCP, MIC10

**Target/Specificity**

CD46 (AAH30594, 36 a.a. ~ 135 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

MCP Antibody (monoclonal) (M07) is for research use only and not for use in diagnostic or therapeutic procedures.


**MCP Antibody (monoclonal) (M07) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **MCP Antibody (monoclonal) (M07) - Images**



Detection limit for recombinant GST tagged CD46 is approximately 1ng/ml as a capture antibody.

### **MCP Antibody (monoclonal) (M07) - Background**

The protein encoded by this gene is a type I membrane protein and is a regulatory part of the complement system. The encoded protein has cofactor activity for inactivation of complement components C3b and C4b by serum factor I, which protects the host cell from damage by complement. In addition, the encoded protein can act as a receptor for the Edmonston strain of measles virus, human herpesvirus-6, and type IV pili of pathogenic *Neisseria*. Finally, the protein encoded by this gene may be involved in the fusion of the spermatozoa with the oocyte during fertilization. Mutations at this locus have been associated with susceptibility to hemolytic uremic syndrome. Alternatively spliced transcript variants encoding different isoforms have been described.

### **MCP Antibody (monoclonal) (M07) - References**

Adenovirus 11p downregulates CD46 early in infection. Gustafsson DJ, et al. *Virology*, 2010 Sep 30. PMID 20638094. Mutations in alternative pathway complement proteins in American patients with atypical hemolytic uremic syndrome. Maga TK, et al. *Hum Mutat*, 2010 Jun. PMID 20513133. Polymorphisms in innate immunity genes and risk of childhood leukemia. Han S, et al. *Hum Immunol*, 2010 Jul. PMID 20438785. Risk of meningioma and common variation in genes related to innate immunity. Rajaraman P, et al. *Cancer Epidemiol Biomarkers Prev*, 2010 May. PMID 20406964. New genetic associations detected in a host response study to hepatitis B vaccine. Davila S, et al. *Genes Immun*, 2010 Apr. PMID 20237496.