

NR2C2 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant NR2C2.****Catalog # AT3100a****Specification**

NR2C2 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC
Primary Accession	P49116
Other Accession	NM_003298
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	65414

NR2C2 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 7182**Other Names**

Nuclear receptor subfamily 2 group C member 2, Orphan nuclear receptor TAK1, Orphan nuclear receptor TR4, Testicular receptor 4, NR2C2, TAK1, TR4

Target/Specificity

NR2C2 (NP_003289, 43 a.a. ~ 152 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

NR2C2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


NR2C2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR2C2 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 KDa) .




Western Blot analysis of NR2C2 expression in transfected 293T cell line by NR2C2 monoclonal antibody (M01), clone 2A5.

Lane 1: NR2C2 transfected lysate (65.414 KDa).

Lane 2: Non-transfected lysate.



Western blot analysis of NR2C2 over-expressed 293 cell line, cotransfected with NR2C2 Validated Chimera RNAi (Cat # AT3100a)



Immunoperoxidase of monoclonal antibody to NR2C2 on formalin-fixed paraffin-embedded human esophagus. [antibody concentration 3 ug/ml]

NR2C2 Antibody (monoclonal) (M01) - Background

Members of the nuclear hormone receptor family, such as NR2C2, act as ligand-activated transcription factors. The proteins have an N-terminal transactivation domain, a central DNA-binding domain with 2 zinc fingers, and a ligand-binding domain at the C terminus. The activated receptor/ligand complex is translocated to the nucleus where it binds to hormone response elements of target genes (Yoshikawa et al., 1996 [PubMed 8661150]).

NR2C2 Antibody (monoclonal) (M01) - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Constitutive association of TGF-beta-activated kinase 1 with the IkappaB kinase complex in the nucleus and cytoplasm of human neutrophils and its impact on downstream processes. Ear T, et al. J Immunol, 2010 Apr 1. PMID 20200282. NF-kappaB activation in T cells requires discrete control of IkappaB kinase alpha/beta (IKKalpha/beta) phosphorylation and IKKgamm ubiquitination by the ADAP adapter protein. Srivastava R, et al. J Biol Chem, 2010 Apr 9. PMID 20164171. Human T cell lymphotropic virus 1 manipulates interferon regulatory signals by controlling the TAK1-IRF3 and IRF4 pathways. Suzuki S, et al. J Biol Chem, 2010 Feb 12. PMID 19955181. Human testicular orphan receptor 4 enhances thyroid hormone receptor signaling. Huang YH, et al. J Cell Physiol, 2010 Feb. PMID

19859911.