

NTRK2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant NTRK2. Catalog # AT3122a

Specification

NTRK2 Antibody (monoclonal) (M01) - Product Information

Application WB **Primary Accession** 016620 BC031835 Other Accession Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 91999

NTRK2 Antibody (monoclonal) (M01) - Additional Information

Gene ID 4915

Other Names

BDNF/NT-3 growth factors receptor, GP145-TrkB, Trk-B, Neurotrophic tyrosine kinase receptor type 2, TrkB tyrosine kinase, Tropomyosin-related kinase B, NTRK2, TRKB

Target/Specificity

NTRK2 (AAH31835, 1 a.a. ~ 477 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

NTRK2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

NTRK2 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot



- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NTRK2 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (78.21 KDa).

NTRK2 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation. Mutations in this gene have been associated with obesity and mood disorders. Alternate transcriptional splice variants encoding different isoforms have been found for this gene.

NTRK2 Antibody (monoclonal) (M01) - References

1.Peptide-mediated targeting of liposomes to TrkB receptor-expressing cells.Ranjan S, Sood R, Dudas J, Glueckert R, Schrott-Fischer A, Roy S, Pyykko I, Kinnunen PK.Int J Nanomedicine. 2012;7:3475-85. Epub 2012 Jul 6.2.Hypoxia-inducible Factor-1 (HIF-1) Is a Transcriptional Activator of the TrkB Neurotrophin Receptor Gene.Martens LK, Kirschner KM, Warnecke C, Scholz H.J Biol Chem. 2007 May 11;282(19):14379-88. Epub 2007 Mar 20.