

PSMB6 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a full length recombinant PSMB6.****Catalog # AT3461a****Specification**

PSMB6 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC, E
Primary Accession	P28072
Other Accession	BC000835
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	25358

PSMB6 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 5694**Other Names**

Proteasome subunit beta type-6, Macropain delta chain, Multicatalytic endopeptidase complex delta chain, Proteasome delta chain, Proteasome subunit Y, PSMB6, LMPY, Y

Target/Specificity

PSMB6 (AAH00835, 1 a.a. ~ 239 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PSMB6 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PSMB6 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PSMB6 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (52.03 KDa) .



Immunoperoxidase of monoclonal antibody to PSMB6 on formalin-fixed paraffin-embedded human lymph node tissue. [antibody concentration 1 ~ 10 ug/ml]



Detection limit for recombinant GST tagged PSMB6 is approximately 3ng/ml as a capture antibody.

PSMB6 Antibody (monoclonal) (M01) - Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit in the proteasome. This catalytic subunit is not present in the immunoproteasome and is replaced by catalytic subunit 1i (proteasome beta 9 subunit).

PSMB6 Antibody (monoclonal) (M01) - References

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. Cell, 2005 Sep 23. PMID 16169070. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Mammalian Cdh1/Fzr mediates its own degradation. Listovsky T, et al. EMBO J, 2004 Apr 7. PMID 15029244. Protein-protein interactions among human 20S proteasome subunits and proteasassemblin. Jayarapu K, et al. Biochem Biophys Res Commun, 2004 Feb 6. PMID 14733938.