

SFPQ Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant SFPQ.****Catalog # AT3842a****Specification**

SFPQ Antibody (monoclonal) (M01) - Product Information

Application	WB, IF, E
Primary Accession	P23246
Other Accession	NM_005066
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	76149

SFPQ Antibody (monoclonal) (M01) - Additional Information**Gene ID** 6421**Other Names**

Splicing factor, proline- and glutamine-rich, 100 kDa DNA-pairing protein, hPOMp100, DNA-binding p52/p100 complex, 100 kDa subunit, Polypyrimidine tract-binding protein-associated-splicing factor, PSF, PTB-associated-splicing factor, SFPQ, PSF

Target/Specificity

SFPQ (NP_005057, 269 a.a. ~ 361 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IF~~1:50~200

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


SFPQ Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

SFPQ Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SFPQ Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to SFPQ on HeLa cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.97 KDa) .



SFPQ monoclonal antibody (M01), clone 3B10. Western Blot analysis of SFPQ expression in HeLa S3 NE (Cat # AT3842a)



Detection limit for recombinant GST tagged SFPQ is approximately 0.3ng/ml as a capture antibody.

SFPQ Antibody (monoclonal) (M01) - References

hnRNP M interacts with PSF and p54(nrb) and co-localizes within defined nuclear structures. Marko M, et al. Exp Cell Res, 2010 Feb 1. PMID 19874820. Identification and characterisation of a novel GHR defect disrupting the polypyrimidine tract and resulting in GH insensitivity. David A, et al. Eur J Endocrinol, 2010 Jan. PMID 19812236. Melanotic Xp11 translocation renal cancer: a case with PSF-TFE3 gene fusion and up-regulation of melanogenetic transcripts. Chang IW, et al. Am J Surg Pathol, 2009 Dec. PMID 19809274. Human PSF binds to RAD51 and modulates its homologous-pairing and strand-exchange activities. Morozumi Y, et al. Nucleic Acids Res, 2009 Jul. PMID 19447914. Interplay between polypyrimidine tract binding protein-associated splicing factor and human myometrial progesterone receptors. Tyson-Capper AJ, et al. J Mol Endocrinol, 2009 Jul. PMID 19339282.