

TIMP2 Antibody (monoclonal) (M01J)

Mouse monoclonal antibody raised against a full length recombinant TIMP2. Catalog # AT4243a

Specification

TIMP2 Antibody (monoclonal) (M01J) - Product Information

Application WB, IHC, IF **Primary Accession** P16035 Other Accession BC052605 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 24399

TIMP2 Antibody (monoclonal) (M01J) - Additional Information

Gene ID 7077

Other Names

Metalloproteinase inhibitor 2, CSC-21K, Tissue inhibitor of metalloproteinases 2, TIMP-2, TIMP2

Target/Specificity

TIMP2 (AAH52605, 27 a.a. \sim 220 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

TIMP2 Antibody (monoclonal) (M01J) is for research use only and not for use in diagnostic or therapeutic procedures.

TIMP2 Antibody (monoclonal) (M01J) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


TIMP2 Antibody (monoclonal) (M01J) - Images



Immunofluorescence of monoclonal antibody to TIMP2 on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (47.08 KDa).



Immunoperoxidase of monoclonal antibody to TIMP2 on formalin-fixed paraffin-embedded human



placenta. [antibody concentration 3 ug/ml]

TIMP2 Antibody (monoclonal) (M01J) - Background

This gene is a member of the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique role among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. As a result, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing the proliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity in tissues undergoing remodelling of the extracellular matrix.

TIMP2 Antibody (monoclonal) (M01J) - References

Clinical Impact of MMP and TIMP Gene Polymorphisms in Gastric Cancer. Alakus H, et al. World J Surg, 2010 Aug 21. PMID 20730428.Matrix metalloproteinase-3 promoter polymorphisms but not dupA-H. pylori correlate to duodenal ulcers in H. pylori-infected females. Yeh YC, et al. BMC Microbiol, 2010 Aug 13. PMID 20707923.Common genetic polymorphisms in Moyamoya and atherosclerotic disease in Europeans. Roder C, et al. Childs Nerv Syst, 2010 Aug 6. PMID 20694560.A genetic association study of maternal and fetal candidate genes that predispose to preterm prelabor rupture of membranes (PROM). Romero R, et al. Am J Obstet Gynecol, 2010 Jul 29. PMID 20673868.Genetic variants in COL2A1, COL11A2, and IRF6 contribute risk to nonsyndromic cleft palate. Nikopensius T, et al. Birth Defects Res A Clin Mol Teratol, 2010 Jul 29. PMID 20672350.