

TP53 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant TP53.****Catalog # AT4310a****Specification**

TP53 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC, IF, IP
Primary Accession	P04637
Other Accession	BC003596
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	43653

TP53 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 7157**Other Names**

Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53, TP53, P53

Target/Specificity

TP53 (AAH03596, 94 a.a. ~ 201 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

IF~~1:50~200

IP~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


TP53 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

TP53 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TP53 Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to TP53 on A-431 cell. [antibody concentration 10 ug/ml]




Immunoprecipitation of TP53 transfected lysate using anti-TP53 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with TP53 MaxPab rabbit polyclonal antibody.



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 KDa) .




TP53 monoclonal antibody (M01), clone 2C3 Western Blot analysis of TP53 expression in A-431 (Cat # AT4310a)




Western Blot analysis of TP53 expression in transfected 293T cell line by TP53 monoclonal antibody (M01), clone 2C3.

Lane 1: TP53 transfected lysate(43.7 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to TP53 on formalin-fixed paraffin-embedded human salivary gland. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged TP53 is approximately 1ng/ml as a capture antibody.

TP53 Antibody (monoclonal) (M01) - Background

This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms, which can regulate p53 transcriptional activity.

TP53 Antibody (monoclonal) (M01) - References

Hepatocellular carcinoma and the underlying mechanisms. Oyagbemi AA, et al. Afr Health Sci, 2010 Mar. PMID 20811532. PTEN tumor suppressor plays less prognostic role than P53 tumor suppressor in diffuse large B-cell lymphoma. Liu YY, et al. Leuk Lymphoma, 2010 Sep. PMID 20807096. BRAF Mutation Is Rare in Advanced-Stage Low-Grade Ovarian Serous Carcinomas. Wong KK, et al. Am J Pathol, 2010 Aug 27. PMID 20802181. Role of p53 and p21 polymorphisms in the risk of cervical cancer among Chinese women. Jiang P, et al. Acta Biochim Biophys Sin (Shanghai), 2010 Sep. PMID 20732856. Association of Glutathione S-Transferase, EPHX, and p53 codon 72 Gene Polymorphisms with Adult Acute Myeloid Leukemia. Chauhan PS, et al. DNA Cell Biol, 2010 Aug 23. PMID 20731606.