

PE-Cy7 Anti-Human CD11b (ICRF44) Antibody
Catalog # ATB10239**Specification****PE-Cy7 Anti-Human CD11b (ICRF44) Antibody - Product Information**

Application	FC
Isotype	Mouse IgG1, kappa
Concentration	5 uL (1 ug)/test
Reactivity	Human
Formulation	10 mM NaH2PO4, 150 mM NaCl, 0.09% NaN3, 0.1% gelatin, pH7.2
Host	Mouse

PE-Cy7 Anti-Human CD11b (ICRF44) Antibody - Additional Information

Gene ID	3684
Gene Name	ITGAM
Alternative Name(s)	
Mac-1, integrin αM, Itgam, CR3	

Format
PE-Cy7

Preparation

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

Application Notes

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 uL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 uL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10e5 to 1x10e8 cells.

Storage Conditions

2-8°C protected from light


PE-Cy7 Anti-Human CD11b (ICRF44) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PE-Cy7 Anti-Human CD11b (ICRF44) Antibody - Images



Human peripheral blood monocytes were stained with 5 μ L (1 μ g) PE-Cy7 Anti-Human CD11b (ATB10239) (solid line) or 1 μ g PE-Cy7 Mouse IgG1 isotype control (dashed line).



Human peripheral blood monocytes were stained with 5 μ L (1 μ g) PE-Cy7 Anti-Human CD11b (ICRF44) manufactured by Tonbo Biosciences (left panel) or eBioscience (right panel).

PE-Cy7 Anti-Human CD11b (ICRF44) Antibody - Background

The ICRF44 antibody reacts with human CD11b, also known as integrin alpha M. This 165-170 kDa cell surface glycoprotein is part of a family of integrin receptors that mediate adhesion between cells (cell-cell) and components of the extracellular matrix, e.g. fibrinogen (cell-matrix). In addition, integrins are active signaling receptors which recruit leukocytes to inflammatory sites and promote cell activation. Complete, functional integrin receptors consist of distinct combinations of integrin chains which are differentially expressed. Integrin alpha M (CD11b) assembles with Integrin beta-2 (CD18) into a receptor known as Macrophage Antigen-1 (Mac-1) or complement receptor type 3 (CR3). This receptor binds and induces intracellular signaling through ICAM-1, ICAM-2, ICAM-3 and ICAM-4 on endothelial cells and can also facilitate removal of iC3b bearing foreign cells. The ICRF44 antibody is widely used as a marker for CD11b expression on macrophages, granulocytes, and subsets of NK cells. It is reported to be cross-reactive with a number of non-human species including Baboon, Chimpanzee, Cynomolgus, Rhesus and Swine.