

FITC Anti-Human CD19 (SJ25C1) Antibody
Catalog # ATB10446**Specification****FITC Anti-Human CD19 (SJ25C1) Antibody - Product Information**

Application	FC
Isotype	Mouse IgG1, kappa
Concentration	5 µL (0.125 µg)/test
Reactivity	Human
Formulation	10 mM NaH ₂ PO ₄ , 150 mM NaCl, 0.09% NaN ₃ , 0.1% gelatin, pH7.2


FITC Anti-Human CD19 (SJ25C1) Antibody - Additional Information


Gene ID	930
Gene Name	CD19
Alternative Name(s)	
Leu-12, B4	

Format
FITC**Storage Conditions**
2-8°C protected from light**FITC Anti-Human CD19 (SJ25C1) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FITC Anti-Human CD19 (SJ25C1) Antibody - Images



Human peripheral blood lymphocytes were stained with 5 μ L (0.125 μ g) FITC Anti-Human CD19 (ATB10446) (solid line) or 0.125 μ g FITC Mouse IgG1 isotype control (dashed line).

FITC Anti-Human CD19 (SJ25C1) Antibody - Background

The SJ25C1 antibody reacts with human CD19, a 95 kDa glycoprotein which acts as a co-receptor, along with CD21 (CR2), CD81 (TAPA-1) and CD225 (Leu13), in support of the functional B cell receptor (BCR). This complex provides antigen-specific recognition and subsequent activation of B cells to proliferate and differentiate into antibody-secreting cells (plasma cells) or memory B cells, which are crucial for secondary antigen encounter. Upon activation and tyrosine phosphorylation, the CD19 molecule can provide an anchor for cytoplasmic signaling proteins such as GRB2, SOS or PLCG2. CD19 is a lineage-differentiation marker, as its expression is detectable at the earliest B cell stages, through development, and is finally lost upon transition to mature plasma cells.

The SJ25C1 antibody is widely used as a phenotypic marker for CD19 expression on B cells, as well as on dendritic cell subsets

FITC Anti-Human CD19 (SJ25C1) Antibody - References

Piatosa B, Birbach M, Siewiera K, Ussowicz M, Kalwak K, Drabko K, Rekawek A, Tkaczyk K, Kurowski PN. 2013. Cytometry Part B. 84B: 179-186. (Flow Cytometry)

Sekiguchi DR, Smith SB, Sutter JA, Goodman NG, Probert K, Louzoun Y, Rogers W, Luning Prak ET. 2011. Cytometry Part B. 80B: 291-299. (Flow Cytometry)

Dörken B, Möller P, Pezzutto A, Schwartz-Albiez R, Moldenhauer G. B-cell antigens: CD19. In: Knapp W, Dörken B, Gilks WR, et al, ed. Leucocyte Typing IV: White Cell Differentiation Antigens. New York, NY: Oxford University Press; 1989:34-36. (Flow Cytometry)

Karahan GE, Eikmans M, Anholts JDH, Class FHJ and Heidt S. 2014. Clin Exp Immunol. 177(1): 333-340. (Flow Cytometry)

Bansal RR, Mackay CR, Moser B and Ebert M. 2012. Eur J Immunol. 42(1): 110-119. (Flow Cytometry)

Jourdan M, Caraux A, Caron G, Robert N, Fiol G, Rème T, Bolloré K, Vendrell JP, Le Gallou S, Mourcin F, De Vos J, Kassambara A, Duperray C, Hose D, Fest T, Tarte K and Klein B. 2011. J Immunol. 187(8): 3931-3941. (Flow Cytometry)

Goval J-J, Greimers R, Boniver J and de Leval L. 2006. J Histochem Cytochem. 54(1): 75-84.
(Immunofluorescence – frozen tissue)