

**PIP4K2A Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5494**

**Specification**

---

**PIP4K2A Antibody (C-term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IF, FC,E    |
| Primary Accession | <a href="#">P48426</a> |
| Reactivity        | Human, Mouse           |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | H=46, 40;M=46;R=46 KDa |
| Isotype           | Rabbit IgG             |
| Antigen Source    | HUMAN                  |

**PIP4K2A Antibody (C-term) - Additional Information**

**Gene ID** 5305

**Antigen Region**  
329-363

**Other Names**

Phosphatidylinositol 5-phosphate 4-kinase type-2 alpha, 1-phosphatidylinositol 5-phosphate 4-kinase 2-alpha, Diphosphoinositide kinase 2-alpha, PIP5KIII, Phosphatidylinositol 5-phosphate 4-kinase type II alpha, PI(5)P 4-kinase type II alpha, PIP4KII-alpha, PtdIns(4)P-5-kinase B isoform, PtdIns(4)P-5-kinase C isoform, PtdIns(5)P-4-kinase isoform 2-alpha, PIP4K2A, PIP5K2, PIP5K2A

**Dilution**

WB~~1:1000  
IHC-P~~1:25  
IF~~1:25  
FC~~1:25

**Target/Specificity**

This PIP4K2A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 329-363 amino acids from the C-terminal region of human PIP4K2A.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PIP4K2A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PIP4K2A Antibody (C-term) - Protein Information**

**Name** PIP4K2A ([HGNC:8997](#))

### Function

Catalyzes the phosphorylation of phosphatidylinositol 5- phosphate (PtdIns5P) on the fourth hydroxyl of the myo-inositol ring, to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P<sub>2</sub>) (PubMed:<a href="http://www.uniprot.org/citations/23326584" target="\_blank">23326584</a>, PubMed:<a href="http://www.uniprot.org/citations/9367159" target="\_blank">9367159</a>). Has both ATP- and GTP-dependent kinase activities (PubMed:<a href="http://www.uniprot.org/citations/26774281" target="\_blank">26774281</a>). May exert its function by regulating the levels of PtdIns5P, which functions in the cytosol by increasing AKT activity and in the nucleus signals through ING2 (PubMed:<a href="http://www.uniprot.org/citations/18364242" target="\_blank">18364242</a>). May regulate the pool of cytosolic PtdIns5P in response to the activation of tyrosine phosphorylation (By similarity). Required for lysosome-peroxisome membrane contacts and intracellular cholesterol transport through modulating peroxisomal PtdIns(4,5)P<sub>2</sub> level (PubMed:<a href="http://www.uniprot.org/citations/29353240" target="\_blank">29353240</a>). In collaboration with PIP4K2B, has a role in mediating autophagy in times of nutrient stress (By similarity). Required for autophagosome-lysosome fusion and the regulation of cellular lipid metabolism (PubMed:<a href="http://www.uniprot.org/citations/31091439" target="\_blank">31091439</a>). May be involved in thrombopoiesis, and the terminal maturation of megakaryocytes and regulation of their size (By similarity). Negatively regulates insulin signaling through a catalytic-independent mechanism (PubMed:<a href="http://www.uniprot.org/citations/31091439" target="\_blank">31091439</a>). PIP4Ks interact with PIP5Ks and suppress PIP5K-mediated PtdIns(4,5)P<sub>2</sub> synthesis and insulin-dependent conversion to PtdIns(3,4,5)P<sub>3</sub> (PubMed:<a href="http://www.uniprot.org/citations/31091439" target="\_blank">31091439</a>).

### Cellular Location

Cell membrane {ECO:0000250|UniProtKB:O70172}. Nucleus. Lysosome {ECO:0000250|UniProtKB:O70172}. Cytoplasm. Photoreceptor inner segment {ECO:0000250|UniProtKB:O70172}. Cell projection, cilium, photoreceptor outer segment {ECO:0000250|UniProtKB:O70172}. Note=May translocate from the cytosol to the cell membrane upon activation of tyrosine phosphorylation. May translocate from the inner to the outer segments of the rod photoreceptor cells in response to light (By similarity) Localization to the nucleus is modulated by the interaction with PIP4K2B. {ECO:0000250|UniProtKB:O70172, ECO:0000269|PubMed:20583997}

### Tissue Location


Expressed ubiquitously, with high levels in the brain. Present in most tissues, except notably skeletal muscle and small intestine.

### PIP4K2A Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### PIP4K2A Antibody (C-term) - Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human Cervical epithelial adenocarcinoma cell line) cells labeling PIP4K2A with AW5494 at 1/25 dilution, followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (1583138) secondary antibody at 1/400 dilution (green). Confocal image showing cytoplasmic staining on HeLa cell line. Cytoplasmic actin is detected with Alexa Fluor® 555 conjugated with Phalloidin (OB16636430) at 1/100 dilution (red).



All lanes : Anti-PIP4K2A Antibody (C-term) at 1:1000 dilution Lane 1: human brain lysates Lane 2: HeLa whole cell lysates Lane 3: K562 whole cell lysates Lane 4: mouse brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 46 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded H. brain section using PIP4K2A Antibody (C-term)(Cat#AW5494). AW5492 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Flow cytometric analysis of Hela cells using PIP4K2A Antibody (C-term)(green, Cat#AW5494) compared to an isotype control of rabbit IgG(blue). AW5492 was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

### PIP4K2A Antibody (C-term) - Background

Catalyzes the phosphorylation of phosphatidylinositol 5- phosphate (PtdIns5P) on the fourth hydroxyl of the myo-inositol ring, to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P<sub>2</sub>). May exert its function by regulating the levels of PtdIns5P, which functions in the cytosol by increasing AKT activity and in the nucleus signals through ING2. May regulate the pool of cytosolic PtdIns5P in response to the activation of tyrosine phosphorylation. May negatively regulate insulin-stimulated glucose uptake by lowering the levels of PtdIns5P. May be involved in thrombopoiesis, and the terminal maturation of megakaryocytes and regulation of their size.

### PIP4K2A Antibody (C-term) - References

- Boronenkov I.V.,et al.J. Biol. Chem. 270:2881-2884(1995).
- Boronenkov I.V.,et al.Submitted (JAN-2000) to the EMBL/GenBank/DDBJ databases.
- Divecha N.,et al.Biochem. J. 309:715-719(1995).
- Deloukas P.,et al.Nature 429:375-381(2004).
- Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

**PIP4K2A Antibody (C-term) - Citations**

- [Distribution and localization of phosphatidylinositol 5-phosphate, 4-kinase alpha and beta in the brain](#)