

P2RX5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5538

Specification

P2RX5 Antibody (C-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q93086
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=47,48,44 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

P2RX5 Antibody (C-term) - Additional Information

Gene ID 5026

Antigen Region
325-353

Other Names

P2X purinoceptor 5, P2X5, ATP receptor, Purinergic receptor, P2RX5, P2X5

Dilution

IHC-P~~1:10~50

WB~~1:1000

Target/Specificity

This P2RX5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 325-353 amino acids from the C-terminal region of human P2RX5.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

P2RX5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

P2RX5 Antibody (C-term) - Protein Information

Name P2RX5 ([HGNC:8536](#))

Function

[Isoform 6]: ATP-gated nonselective transmembrane cation channel permeable to potassium, sodium and calcium (PubMed:

target="_blank">12761352). Unlike, other P2RX receptors, the P2X5 receptor is also permeable to chloride (PubMed:12761352). May play a supporting role in the inflammatory response (By similarity).

Cellular Location

[Isoform 6]: Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P56373}. Note=The absence of the exon 10-encoded amino acids substantially results in subunit aggregation and retention in the cell cytoplasm

Tissue Location


Expressed at high levels in brain and immune system

P2RX5 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

P2RX5 Antibody (C-term) - Images



All lanes : Anti-P2RX5 Antibody (C-term) at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



P2RX5 Antibody (C-term) (AW5538) immunohistochemistry analysis in formalin fixed and paraffin embedded human cervix tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of P2RX5 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

P2RX5 Antibody (C-term) - Background

The product of this gene belongs to the family of purinoceptors for ATP. This receptor functions as a ligand-gated ion channel. Several characteristic motifs of ATP-gated channels are present in its primary structure, but, unlike other members of the purinoceptors family, this receptor has only a single transmembrane domain. Three transcript variants encoding distinct isoforms have been identified for this gene.

P2RX5 Antibody (C-term) - References

Kotnis, S., et al. Mol. Pharmacol. 77(6):953-960(2010)
Overes, I.M., et al. Cancer Immunol. Immunother. 58(3):429-439(2009)
Dubyak, G.R. Mol. Pharmacol. 72(6):1402-1405(2007)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Duckwitz, W., et al. J. Biol. Chem. 281(51):39561-39572(2006)