

SAMHD1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5632

Specification

SAMHD1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9Y3Z3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=72,69,64,68 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

SAMHD1 Antibody (Center) - Additional Information

Gene ID 25939

Antigen Region
204-230

Other Names

Deoxynucleoside triphosphate triphosphohydrolase SAMHD1, dNTPase, 315-, Dendritic cell-derived IFNG-induced protein, DCIP, Monocyte protein 5, MOP-5, SAM domain and HD domain-containing protein 1, SAMHD1, MOP5

Dilution
WB~~0.25

Target/Specificity

This SAMHD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 204-230 amino acids from the Central region of human SAMHD1.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SAMHD1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SAMHD1 Antibody (Center) - Protein Information

Name SAMHD1 ([HGNC:15925](#))

Function

Protein that acts both as a host restriction factor involved in defense response to virus and as a

regulator of DNA end resection at stalled replication forks (PubMed:19525956, PubMed:21613998, PubMed:21720370, PubMed:22056990, PubMed:23601106, PubMed:23602554, PubMed:24336198, PubMed:26294762, PubMed:26431200, PubMed:28229507, PubMed:28834754, PubMed:29670289). Has deoxynucleoside triphosphate (dNTPase) activity, which is required to restrict infection by viruses, such as HIV-1: dNTPase activity reduces cellular dNTP levels to levels too low for retroviral reverse transcription to occur, blocking early- stage virus replication in dendritic and other myeloid cells (PubMed:19525956, PubMed:21613998, PubMed:21720370, PubMed:22056990, PubMed:23364794, PubMed:23601106, PubMed:23602554, PubMed:24336198, PubMed:25038827, PubMed:26101257, PubMed:26294762, PubMed:26431200, PubMed:28229507). Likewise, suppresses LINE-1 retrotransposon activity (PubMed:24035396, PubMed:24217394, PubMed:29610582). Not able to restrict infection by HIV-2 virus; because restriction activity is counteracted by HIV-2 viral protein Vpx (PubMed:21613998, PubMed:21720370). In addition to virus restriction, dNTPase activity acts as a regulator of DNA precursor pools by regulating dNTP pools (PubMed:23858451). Phosphorylation at Thr-592 acts as a switch to control dNTPase-dependent and -independent functions: it inhibits dNTPase activity and ability to restrict infection by viruses, while it promotes DNA end resection at stalled replication forks (PubMed:23601106, PubMed:23602554, PubMed:29610582, PubMed:29670289). Functions during S phase at stalled DNA replication forks to promote the resection of gapped or reversed forks: acts by stimulating the exonuclease activity of MRE11, activating the ATR-CHK1 pathway and allowing the forks to restart replication (PubMed:29670289). Its ability to promote degradation of nascent DNA at stalled replication forks is required to prevent induction of type I interferons, thereby preventing chronic inflammation (PubMed:27477283, PubMed:29670289). Ability to promote DNA end resection at stalled replication forks is independent of dNTPase activity (PubMed:29670289). Enhances immunoglobulin hypermutation in B-lymphocytes by promoting transversion mutation

(By similarity).

Cellular Location

Nucleus. Chromosome Note=Localizes to sites of DNA double-strand breaks in response to DNA damage.

Tissue Location


Expressed in heart, skeletal muscle, spleen, liver, small intestine, placenta, lung and peripheral blood leukocytes (PubMed:11064105). No expression is seen in brain and thymus (PubMed:11064105).

SAMHD1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SAMHD1 Antibody (Center) - Images



All lanes : Anti-SAMHD1 Antibody (Center) at 1:2000 dilution Lane 1: THP-1 whole cell lysate Lane 2: Ramos whole cell lysate Lane 3: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 72 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

SAMHD1 Antibody (Center) - Background

This gene may play a role in regulation of the innate immune response. The encoded protein is upregulated in response to viral infection and may be involved in mediation of tumor necrosis factor-alpha proinflammatory responses. Mutations in this gene have

been associated with Aicardi-Goutieres syndrome. [provided by RefSeq].

SAMHD1 Antibody (Center) - References

Tomkova, H., et al. Eur J Dermatol 20(3):411-413(2010)
Dale, R.C., et al. Am. J. Med. Genet. A 152A (4), 938-942 (2010) :
Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Rice, G.I., et al. Nat. Genet. 41(7):829-832(2009)
Liao, W., et al. Proteomics 8(13):2640-2650(2008)