10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Zebrafish papl Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb18704c

Specification

Zebrafish papl Antibody (C-term) - Product Information

Application	**WB,E**
Primary Accession	A5D6U8
Reactivity	**Zebrafish**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**378-413**

Zebrafish papl Antibody (C-term) - Additional Information

Gene ID 571830

Other Names
Iron/zinc purple acid phosphatase-like protein, papl

Target/Specificity
This Zebrafish papl antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 378-413 amino acids of zebrafish papl.

Dilution
WB~~1:1000
E~~Use at an assay dependent concentration.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
Zebrafish papl Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Zebrafish papl Antibody (C-term) - Protein Information

Name acp7 {ECO:0000250|UniProtKB:Q6ZNF0}

Cellular Location
Secreted.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Zebrafish papl Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Zebrafish papl Antibody (C-term) - Images



Western blot analysis of lysates from zebrafish brain, ZF4 tissue lysate (from left to right), using Zebrafish papl Antibody (C-term) (Cat. #Azb18704c). Azb18704c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.