

(DANRE) chst1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb18710c

Specification

(DANRE) chst1 Antibody (C-term) - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Antigen Region	WB,E <u>O6DBY9</u> Zebrafish Rabbit Polyclonal Rabbit IgG
Antigen Region	341-375

(DANRE) chst1 Antibody (C-term) - Additional Information

Gene ID 445124

Other Names Carbohydrate sulfotransferase 1, Keratan sulfate Gal-6 sulfotransferase, KS6ST, KSGal6ST, KSST, chst1

Target/Specificity This (DANRE) chst1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 341-375 amino acids of DANRE chst1.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions (DANRE) chst1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) chst1 Antibody (C-term) - Protein Information

Name chst1

Function Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of galactose (Gal) residues of keratan.



Cellular Location


Golgi apparatus membrane; Single- pass type II membrane protein

(DANRE) chst1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

(DANRE) chst1 Antibody (C-term) - Images



Western blot analysis of lysates from zebra fish muscle, Zebrafish tissue lysate (from left to right), using (DANRE) chst1 Antibody (C-term)(Cat. #Azb18710c). Azb18710c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

(DANRE) chst1 Antibody (C-term) - Background

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of galactose (Gal) residues of keratan (By similarity).