

Zebrafish mycl1a Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb18711a**Specification**

Zebrafish mycl1a Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9PSI9
Reactivity	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	80-113

Zebrafish mycl1a Antibody (N-term) - Additional Information**Gene ID** 405873**Other Names**

Protein L-Myc-1a, Protein L-Myc 1, zL-Myc, mycl1a {ECO:0000312|EMBL:AAH677231}, lmyc1, mycl1

Target/Specificity

This Zebrafish mycl1a antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 80-113 amino acids from the N-terminal region of zebrafish mycl1a.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Zebrafish mycl1a Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Zebrafish mycl1a Antibody (N-term) - Protein Information**Name** mycl1a {ECO:0000312|EMBL:AAH67723.1}**Synonyms** lmyc1, mycl1

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P12524, ECO:0000255|PROSITE-ProRule:PRU00981}


Tissue Location

Uterus..

Zebrafish mycl1a Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zebrafish mycl1a Antibody (N-term) - Images

Western blot analysis of lysates from zebrafish muscle, zebrafish tissue lysate (from left to right), using Zebrafish mycl1a Antibody (N-term) (Cat. #Azb18711a). Azb18711a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Zebrafish mycl1a Antibody (N-term) - References

Schreiber-Agus N.,et al.Mol. Cell. Biol. 13:2765-2775(1993).