

Animal-Free Recombinant Human SDF-1ß (CXCL12)

Catalog # PBG10547

Specification

Animal-Free Recombinant Human SDF-1ß (CXCL12) - Product Information

Animal-Free Recombinant Human SDF-1β (CXCL12) - Additional Information

Description

SDF- 1α and β are stromal derived CXC chemokines, and signal through the CXCR4 receptor. SDF- 1α and β chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV suppressive activity in cells expressing the CXCR4 receptor. Recombinant human SDF- 1β is an 8.5 kDa protein containing 72 amino acid residues.

BiologicalActivity

Determined by its ability to chemoattract human peripheral T cells activated with PHA and IL-2 using a concentration range of 20-80 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is $<0.1 \text{ ng}/\mu\text{g}$ of protein ($<1\text{EU}/\mu\text{g}$).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Animal-Free Recombinant Human SDF-1 β (CXCL12) is for research use only and not for use in diagnostic or therapeutic procedures.

Animal-Free Recombinant Human SDF-1ß (CXCL12) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Animal-Free Recombinant Human SDF-1β (CXCL12) - Images



